

JHH- Documenting Patient Controlled Analgesia (PCA)

Required PCA documentation

When an Emergency Department patient requires a PCA, there is specific documentation to complete.



Try It Out

1. Document the administration of the ordered medication using the MAR toolbox in the ED Narrator or the MAR activity:

- PCA medication administration will require a dual signoff by a second RN

2. To document an ongoing assessment of the PCA, open the **ED Narrator** (1), choose **Assessments** (2), click the **Add** button (3), select the **PCA assessment** (4), and then click **Accept** (5):

3. This flowsheet allows you to document specifics surrounding the patient's PCA use. Once finished, click **Accept**:

Event Log | Patient Summary | Physical Diagram | Orders

Show: Deleted Status Changes Flowsheets/Assessments LDA Notes Tx Trm Orders Med Admin Flowsheet File

▷ morphine PCA 250 mg/50 mL NS (5 mg/mL)

▽ HYDRomorphone (DILAUDID) PCA 20mg/100mL NS (0.2 mg/mL)

Continuous Infusion Rate (mg/hr)	<input type="text" value="1"/>	PCA dose (mg)	<input type="text" value="0.2"/>
Lockout Interval (min)	<input type="text" value="10"/>	Max Doses/Hour	<input type="text" value="4"/>
The maximum allowable doses that the patient may receive per hour from pushing the remote dose button.			
Reservoir Volume (mL)	<input type="text" value="8"/>	Total Amount Given (mg)	<input type="text" value="0.6"/>
The total volume in mL remaining in the cassette/syringe/bag/reservoir at the time of the reading.			
Number of PCA Doses Given	<input type="text" value="3"/>	Number of Attempts	<input type="text" value="4"/>
Loading Dose (mg)	<input type="text"/>	Clinician Bolus Dose (mg)	<input type="text"/>
Volume Infused (mL)	<input type="text"/>		
Amount of volume infused since last documentation.			

▷ Sensation/Dermatomes

▷ Sedation Scale

▷ Motor Assessment

▷ Vitals

▷ Pain Assessment

Accept Cancel